



Life Threatening Differential Chest Pain Diagnoses Supplemental Guide

The North Dakota Department of Health is not responsible for inaccuracies contained herein. No responsibility is assumed for damages or liabilities arising from accuracy, content error, or omission.

Aortic Aneurysm (Ruptured)

Characteristic symptoms:

- Appears pale and unwell, cold, sweaty and hypotensive
- Tachycardia, weak and thready pulse
- There may be a pulsatile mass in the vicinity of the bifurcation of the aorta. This is a few centimeters above the umbilicus and a little to the left.
- Aneurysm may be tender and a bruit may be audible. Bleeding causes peritoneal irritation and it may appear as an acute abdomen.
- Occasionally atypical presentation will occur . ex. intestinal obstruction from hematoma or an apparent irreducible inguinal hernia.

Diagnosis:

Obtain chest x-ray: Thoracic aortic aneurysm may cause an enlarged base of aorta.

Abdominal X-ray: will reveal abdominal aortic aneurysm lesion in about 75% of cases

Portable ultrasound

Other investigations: CT angiography will confirm the diagnosis, but is time consuming, so transferring patient to tertiary center may be preferred

Treatment:

Emergency surgical or endovascular repair

Aortic Dissection

Characteristic symptoms:

- Sudden severe chest or upper back pain, often described as a tearing, ripping or shearing sensation, that radiates to the neck or down the back
- Loss of consciousness



- Shortness of breath
- Sudden difficulty speaking, loss of vision, weakness or hemiplegia
- Weak pulse/decreased BP in one arm compared with the other

Diagnosis:

Obtain chest x-ray- look for widened mediastinum or aortic knob

CTA, MRI, and TEE

Treatment:

Emergency surgical intervention

Pericardial Tamponade

Characteristic symptoms:

- Chest pain radiating to neck, shoulders, or back
- Hypotension and weak pulse
- Distended neck veins
- Tachycardia combined with muffled heart sounds due to the expanding layer of fluid inside the pericardium
- Blood pressure that falls (pulsus paradoxical) when breathing deeply
- Discomfort that's relieved by sitting or leaning forward
- Trouble breathing or taking deep breaths
- Anxiety and restlessness
- Weakness
- Tachypnea
- Syncope
- Weak or absent peripheral pulses

Diagnosis:

Chest x-ray may show enlarged globular-shaped heart

EKG may show low voltage QRS complexes



Echocardiography is diagnostic test of choice

CTA or MRI

Treatment:

Pericardiocentesis

Tension Pneumothorax

Characteristic symptoms:

- Sudden onset of chest pain that is sharp in nature and may lead to feelings of tightness in the chest.
- Shortness of breath
- Hypotension, tracheal deviation and neck vein distention
- Tachycardia
- Tachypnea
- Cough
- Fatigue
- Cyanosis

Diagnosis:

Decreased or absent breath sounds over the affected lung. Affected hemithorax is also hyperresonant to percussion and often feels distended, tense and poorly compressible to palpation.

Chest X-ray

Treatment:

A small pneumothorax without underlying lung disease may resolve on its own in one to two weeks. A larger pneumothorax and a pneumothorax associated with underlying lung disease often require aspiration of the free air and/or placement of a chest tube to evacuate the air.

Pulmonary Embolism

Characteristic symptoms:

- Shortness of breath that may occur suddenly
- Sudden, sharp chest pain that may become worse with deep breathing or coughing
- Calf pain/swelling



- Rapid heart rate
- Rapid breathing
- Sweating
- Anxiety
- Coughing up blood or pink, foamy mucus
- Syncope
- Heart palpitations

Diagnosis:

Evaluate Wells Score (see below)

Obtain D-Dimer- high levels may be diagnostic of PE

Ultrasound, CT, pulmonary angiogram, MRI

Wells Criteria for Pulmonary Embolism	
Clinical Feature	Points
Clinical s/s of DVT (<i>minimum of leg swelling and pain with palpitation of deep veins</i>)	3
An alternative diagnosis is less likely than PE	3
Heart rate >100 beats per minute	1.5
Immobilization \geq 3 days or surgery in the previous 4 weeks	1.5
Hemoptysis	1
Malignancy (on treatment/treated in the past 6 months/palliative)	1
Clinical Probability	
PE likely	>4
PE unlikely	\leq 4

Treatment:

Administration of anticoagulants and thrombolytics

Endovascular clot removal

Placement of vena cava filter