

ANOGENITAL LESIONS – HUMAN PAPILLOMAVIRUS - RD 7

DEFINITION

External HPV lesions are warty growths which can be found on the penis, vulva, perineum, vagina, cervix, urethra, and perianal areas. 90% of anogenital warts are caused by non-oncogenic HPV types 6 or 11. Oncogenic, high-risk HPV infection (e.g., HPV types 16 and 18) causes the majority of cervical, penile, vulvar, vaginal, anal, and oropharyngeal cancers and precancers, Persistent oncogenic HPV infection is the strongest risk factor for the development of HPV-attributable precancers and cancers, however, the only HPV-associated cancer for which routine screening is recommended is cervical cancer.

SUBJECTIVE

May include:

1. HPV lesions on or near the genital area
2. Pruritic and/or postcoital burning sensation or spotting
3. Known contact with an HPV-infected person
4. No symptoms

OBJECTIVE

Diagnosis is typically made from clinical presentation.

May include:

1. Small to large, dry wart-like growths on or near the genital area or oral areas
2. Single or multiple soft, fleshy, papillary keratinized growths
3. Acetopositive lesions on or near the genital area. (Acetic acid application is not a specific test for HPV infections and is not generally recommended)

LABORATORY

May include:

1. Vaginitis/cervicitis screening, as appropriate
2. STI testing as indicated
3. Pap smear with or without HPV testing, as age appropriate
4. Syphilis testing

*HPV typing is not recommended for anogenital wart diagnosis; it is not confirmatory and does not guide treatment.

ASSESSMENT

Anogenital HPV

PLAN

1. If left untreated, anogenital warts can resolve spontaneously, remain unchanged or increase in size or number. Because warts might spontaneously resolve within 1 year, an acceptable alternative for some persons is to forego treatment and wait for spontaneous resolution.
2. Recommended patient applied treatment regimens for external HPV:
 - a. Imiquimod 3.75% or 5% cream. Imiquimod 5% apply thin layer of cream to lesions 3 times a week at bedtime; rub the cream until it vanishes. Wash hands after application. Leave on for 6-10 hours and then wash the area with mild soap and water. Continue treatment until HPV lesions are resolved for up to 16 weeks. Imiquimod 3.75% is applied the same way as above but is used nightly for < 8 weeks. Not recommended in pregnancy. OR

- b. Podofilox 0.5% solution or gel. Apply 2 times/day for 3 days, then no treatment for 4 days. May repeat for 4 cycles. Wash hands after application. Not recommended in pregnancy. OR
 - c. Sinecatechins 15% ointment. Apply to HPV lesions 3 times/day for up to 16 weeks. Wash hands after application. Not approved for use in pregnancy. May weaken condoms and diaphragms.
3. Recommended Provider applied therapies:
- a. Cryotherapy with liquid nitrogen. Apply liquid nitrogen to lesion and base of lesion for a few seconds, allow thawing then repeat freeze x1 or x2. May reapply treatment every 7-14 days. Not recommended for internal vaginal warts. OR
 - b. Trichloroacetic acid (TCA 80-90%) solution (only available through compounding pharmacy) or bichloroacetic acid (BCA 80-90%) solution. Apply a small amount to the HPV lesions and allow to dry. Avoid contact with surrounding skin. Reapply every 1-2 weeks. May be used on vaginal/introital HPV lesions as well. OR
 - c. Surgical removal by tangential scissor excision, tangential shave excision, curettage, laser, or electro-surgery.
4. Most anogenital warts respond within 3 months of therapy. A new treatment modality should be selected if there is no significant response after a complete course of treatment or if there are therapy-related side effects.

CLIENT EDUCATION

1. Provide client education handout(s). Review the manufacturer's instructions. Review symptoms, treatment options, and medication side effects
2. Advise the client to avoid intercourse during treatment. Imiquimod may weaken latex condoms and diaphragms
3. Instruct on genital self-exam
4. Recommend client RTC prn for treatment, annually, and prn problems
5. Advise the client on available HPV vaccinations
6. Explain that treatment does not eradicate the HPV virus and that HPV lesions may reoccur
7. Provide education on safer sex practices

CONSULT / REFER TO PHYSICIAN

1. Vaginal wall lesions or visible lesions on the cervix
2. Palpable or suspected rectal wall lesions
3. Lesions non-responsive to treatment
4. Immunocompromised patient
5. Surgical removal for extensive HPV lesions
6. Biopsy to confirm or determine diagnosis of lesion if diagnosis is uncertain
7. Refer or consult if pregnant

REFERENCES

1. Marrazzo, J. & Park, I. (2025). Reproductive tract infections, including HIV and other sexually transmitted infections. In R. A. Hatcher, P. Cason, C. Cwiak, A. Edelman, D. Kowal, J. M. Marrazzo,., ... M.S. Policar (Eds.). *Contraceptive technology* (pp. 621- 666). (22nd ed.). Jones & Bartlett Learning.
2. U.S. Centers for Disease and Control. (2021). Sexually transmitted infections treatment guidelines, 2021. *Morbidity and Mortality Weekly Report*, 70(4); 100-106. Retrieved from <https://www.cdc.gov/std/treatment-guidelines/STI-Guidelines-2021.pdf>
3. U.S. Centers for Disease and Control. (2024). *About genital HPV infection*. Retrieved from https://www.cdc.gov/sti/about/about-genital-hpv-infection.html?CDC_AAref_Val=https://www.cdc.gov/STD/HPV/STDFact-HPV.htm