

Immunizations: 18 Months - 18 Years

Immunizations are listed under the earliest recommended age

Influenza and COVID-19 vaccines are recommended annually.

18 mos

Hep B - 3rd Dose
DTaP - 4th Dose
Poliovirus - 3rd Dose
*Hep A - 2nd Dose

4 - 6 years

DTaP - 5th Dose
Poliovirus - 4th Dose
MMR - 2nd Dose
VAR - 2nd Dose

7 - 10 years

*HPV

11 - 12 years

Tdap - 1 Dose
*HPV
Meningococcal - 1st Dose

16 years

Meningococcal - 2nd Dose
Meningococcal B - 1st Dose

17 - 18 years

Meningococcal B - 2nd Dose

In the event of a pregnancy, an RSV immunization is recommended during pregnancy at least 14 days before delivery.

You may require additional vaccines if you are at an increased risk for certain diseases. Talk to your health care provider about what vaccines are right for you.

*Hep A 2-dose series with a minimum of 6 months between doses between ages 12-23 months

Meningococcal - minimum age of 2 months, routine vaccination at 11-12 Years (2-dose series) and 16 years

*HPV - Routinely recommended at 11-12 years, but can start at 9 years; 2- or 3-dose series depending on age at initial vaccination

*Influenza vaccination recommended annually

*COVID-19 vaccine recommended annually

Source: CDC

Immunizations: Key

Listed As	Associated Disease	Associated Immunization/Vaccine
RSV	<i>Respiratory syncytial virus (RSV)</i>	RSV-mAb (Nirsevimab) immunization
Hep B	<i>Hepatitis B</i>	HepB
DTap	<i>Diphtheria, tetanus & acellular pertussis</i>	DTaP (<7 years)
Poliovirus	<i>Inactivated poliovirus</i>	IPV (<18 years)
Hep A	<i>Hepatitis A</i>	HepA
Meningococcal	<i>Meningococcal</i>	MENACWY-CRM (≥2mos), MenACWYT-TT (≥2 years)
MMR	<i>Measles, mumps & rubella</i>	MMR
VAR	<i>Varicella</i>	VAR
Tdap	<i>Tetanus, diphtheria & acellular pertussis</i>	Tdap (≥7 years)
HPV	<i>Human papillomavirus</i>	HPV
Meningococcal B	<i>Meningococcal B</i>	MENB-4C, MenB-FHbp
Influenza	<i>Influenza (Flu)</i>	IIV4 or LAIV4
COVID-19	<i>COVID-19</i>	1vCOV-mRNA, 1vCOV-aPS