

Immunizations: Birth - 15 Months

Immunizations are listed under the earliest recommended age

Influenza and COVID-19 vaccines are recommended annually.

Birth

*RSV
Hep B - 1st Dose

1 mo

Hep B - 2nd Dose

2 mos

Rotavirus - 1st Dose
DTaP - 1st Dose
Hib - 1st Dose
PCV15, PCV20 - 1st Dose
Poliovirus - 1st Dose

4 mos

Rotavirus - 2nd Dose
DTaP - 2nd Dose
Hib - 2nd Dose
PCV15, PCV20 - 2nd Dose
Poliovirus - 2nd Dose

6 mos

Hep B - 3rd Dose
*Rotavirus - 3rd Dose
DTaP - 3rd Dose
*Hib - 3rd Dose
PCV - 3rd Dose
Poliovirus - 3rd Dose

12 mos

Hep B - 3rd Dose
*Hib - 3rd/4th Dose
PCV/PCV20 - 4th Dose
MMR - 1st Dose
VAR - 1st Dose
*Hep A - 1st Dose

15 mos

DTaP - 4th Dose

*RSV - RSV immunization at birth if mother did not receive RSV vaccine at least 14 days prior to delivery

*Rotavirus - 2-dose series and 3-dose series options

*Hib - 3-dose series and 4-dose series options

*Hep A 2-dose series with a minimum of 6 months between doses between ages 12-23 months

You may require additional vaccines if you are at an increased risk for certain diseases. Talk to your health care provider about what vaccines are right for you.

Immunizations: Key

Listed As	Associated Disease	Associated Immunization/Vaccine
RSV	<i>Respiratory syncytial virus (RSV)</i>	RSV-mAb (Nirsevimab) immunization
Hep B	<i>Hepatitis B</i>	HepB
Rotavirus	<i>Rotavirus (RV)</i>	RV1 (2-dose series) or RV5 (3-dose series)
DTaP	<i>Diphtheria, tetanus & acellular pertussis</i>	DTaP (<7 years)
Hib	<i>Haemophilus influenzae type b</i>	Hib
PCV	<i>Pneumococcal conjugate</i>	PCV15, PCV20
Poliovirus	<i>Inactivated poliovirus</i>	IPV (<18 years)
COVID-19	<i>COVID-19</i>	1vCOV-mRNA, 1vCOV-aPS
Influenza	<i>Influenza (Flu)</i>	IIV3 or LAIV3
MMR	<i>Measles, mumps & rubella</i>	MMR
VAR	<i>Varicella</i>	VAR
Hep A	<i>Hepatitis A</i>	HepA
Meningococcal	<i>Meningococcal</i>	MENACWY-CRM (≥2mos), MenACWYT-TT (≥2 years)